COVID UPDATE

Robert Peltier, MD Chief Medical Officer North Oaks Health System





UNITED STATES COVID-19 STATISTICS AS OF JANUARY 8, 2022

	DAILY AVG. ON JAN. 8	14-DAY CHANGE	TOTAL REPORTED
Cases	656,478	+226%	59,848,908
Tests	1,765,601	+15%	
Hospitalized	127,225	+78%	
Deaths	1,524	+12%	835,835

Deaths










All time

Last 90 days




Daily average: 9,967

New cases: 14,802




DAILY CASES IN LOUISIANA (PREDICTIVE MODEL)



SOURCE: IHME WEBSITE




DAILY DEATHS IN LOUISIANA (PREDICTIVE MODEL)



SOURCE: IHME MODEL




HOSPITAL RESOURCE USE IN LOUISIANA




SOURCE: IHME WEBSITE

All begs needed === ICU begs neede





VACCINATION METRICS BY AGE IN LOUISIANA



Do I qualify for a COVID-19 vaccine booster and which one?

Which primary vaccine series did you complete?	Pfizer-BioNTech	Moderna	Janssen (J&J)
You can get a booster if:	It's been at least 5 months since completing a primary series AND you are:	It's been at least 5 months since completing a primary series AND you are:	It's been at least 2 months since completing primary vaccination AND you are:
	Age 12+	Age 18+	Age 18+
If eligible, you can get a booster of:	Pfizer-BioNTech* Moderna Janssen (J&J) *Only Pfizer-BioNTech can be used as a booster in those age 12-17.	Moderna Pfizer-BioNTech Janssen (J&J)	Janssen (J&J) Pfizer-BioNTech Moderna

CDC ISOLATION GUIDELINES



Updated CDC Isolation and Quarantine Recommendations

If You Test **POSITIVE** for COVID-19 (regardless of vaccination status)

If You were **EXPOSED** to COVID-19 & Boosted

If You were **EXPOSED** to COVID-19 & Unvaccinated OR Vaccinated >6 mo. ago with Pfizer or Moderna vaccine or >2 mo. ago with J&J vaccine





Stay home Test if Continue to Possible wear a mask around others

If you can't quarantine you must wear a mask for 10 days.

If you develop symptoms get a test and stay home.

cdc.gov/coronavirus

CURRENT THERAPIES EFFECTIVE AGAINST THE OMICRON VARIANT





NDC 0173-0901-86 Rx Only



Sotrovimab injection 500 mg/8 mL (62.5 mg/mL)

For intravenous infusion after further dilution.

Contains One 8-mL Single-Dose Vial.

Discard Unused Portion.

For Use Under Emergency Use Authorization (EUA)



ORAL ANTIVIRAL COVID TREATMENTS







	Molnupiravir (Merck)	Paxlovid (Pfizer)
Efficacy in high-risk patients, reduction of hospitalizations/deaths at 28 days	30% 9.7 vs 6.8%	89% 8.2 vs 0.7%
Deaths in placebo vs drug	9 vs 0	7 vs 0
Duration of therapy (twice daily)	5 days	5 days
Given with co-drug to promote half- life	No	Yes, ritonavir
Drug interactions (CYP3A4)	Minimal	Yes, many statins, blood thinners
Repurposed	Yes, Equine encephalitis Planned to test for RSV, influenza, redirected	No, Covid specific New chemical entity adapted from an anti-SARS molecule
Mechanism	Nucleoside analog; Induces mutations	Inhibits Mpro, not mutagenic
Active against all variants tested	Yes	Yes
Cost	~\$700	~\$500

QUESTIONS?

